

(Tel: 400-999-8863 ■ Email:Upingbio.163.com



DNJB2 Polyclonal Antibody

YP-Ab-06603
IgG
Human;Rat;Mouse;
WB;ELISA
DNAJB2 HSJ1 HSPF3
DnaJ homolog subfamily B member 2 (DnaJ protein homolog 1) (Heat shock 40 kDa protein 3) (Heat shock protein J1) (HSJ-1)
Synthesized peptide derived from human protein . at AA range: 260-340
DNJB2 Polyclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
35kD
[Isoform 2]: Cytoplasm . Nucleus .; [Isoform 1]: Endoplasmic reticulum membrane; Lipid-anchor; Cytoplasmic side .
More abundantly expressed in neocortex, cerebellum, spinal cord and retina where it is expressed by neuronal cells (at protein level) (PubMed:1599432, PubMed:12754272). Detected at much lower level in non-neuronal tissues including kidney, lung, heart, skeletal muscle, spleen and testis (at protein level) (PubMed:12754272, PubMed:1599432). Isoform 1 is more abundant in neocortex and cerebellum compared to isoform 2 (at protein level) (PubMed:12754272).
alternative products:Experimental confirmation may be lacking for some isoforms,similarity:Contains 1 J domain.,similarity:Contains 2 UIM (ubiquitin-interacting motif) repeats.,tissue specificity:Brain (neuronal layers). Weakly, in skeletal muscle and spleen.,
rreally, in exercial masers and spissing



UpingBio technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from PC12 cells, primary antibody was diluted at 1:1000, 4°over night