


## XCL2 Polyclonal Antibody

Catalog No	YP-Ab-06662
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	XCL2 SCYC2
Protein Name	Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 1-50
Specificity	XCL2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	12kD
Cell Pathway	
och i diliway	Secreted.
Tissue Specificity	Secreted. Blood,Placenta,
Tissue Specificity	Blood,Placenta,  function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine
Tissue Specificity Function	Blood,Placenta,  function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family.,  function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine
Tissue Specificity Function Background matters needing	Blood,Placenta,  function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family.,  function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family.,



## **Products Images**



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night