







## CDK18 Polyclonal Antibody

Catalog No	YP-Ab-06741
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	CDK18 PCTK3
Protein Name	Cyclin-dependent kinase 18 (EC 2.7.11.22) (Cell division protein kinase 18) (PCTAIRE-motif protein kinase 3) (Serine/threonine-protein kinase PCTAIRE-3)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	CDK18 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	<b>,</b>
Synonyms Observed Band	51kD
-	, and the second
Observed Band	, and the second
Observed Band Cell Pathway	51kD  Isoform 2 expression is limited to several subcortical nuclei of the basal gangli and
Observed Band Cell Pathway Tissue Specificity	51kD  Isoform 2 expression is limited to several subcortical nuclei of the basal gangli and the spinal cord. Isoform 1 is widely expressed.
Observed Band Cell Pathway Tissue Specificity Function	51kD  Isoform 2 expression is limited to several subcortical nuclei of the basal gangli and the spinal cord. Isoform 1 is widely expressed. similarity:Belongs to the protein kinase superfamily.,
Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Isoform 2 expression is limited to several subcortical nuclei of the basal gangli and the spinal cord. Isoform 1 is widely expressed. similarity:Belongs to the protein kinase superfamily., similarity:Belongs to the protein kinase superfamily.,



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com

Website: www.upingBio.com	

Products Images