






BECP1 Polyclonal Antibody

Catalog No	YP-Ab-06788
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	BECN1P1
Protein Name	Putative beclin-1-like protein (Beclin-1 autophagy-related pseudogene 1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	BECP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
	> 000/
Purity	≥90%
Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 47kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 47kD Cytoplasm .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 47kD Cytoplasm .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year 47kD Cytoplasm. Present in fetal and adult brain (at protein level). BECN2 (Beclin 2) is a Protein Coding gene. Among its related pathways are Autophagy Pathway and Kaposi's sarcoma-associated herpesvirus infection. An important paralog of this gene is BECN1. Involved in 2 distinct lysosomal degradation pathways: acts as a regulator of autophagy and as a regulator of G-protein coupled receptors turnover. Regulates degradation in lysosomes of a
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background	-20°C/1 year 47kD Cytoplasm. Present in fetal and adult brain (at protein level). BECN2 (Beclin 2) is a Protein Coding gene. Among its related pathways are Autophagy Pathway and Kaposi's sarcoma-associated herpesvirus infection. An important paralog of this gene is BECN1. Involved in 2 distinct lysosomal degradation pathways: acts as a regulator of autophagy and as a regulator of G-protein coupled receptors turnover. Regulates degradation in lysosomes of a variety of G-protein coupled receptors via its interaction with GPRASP1/GASP1.



Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night