







## LAMB4 Polyclonal Antibody

Catalog No	YP-Ab-06820
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	LAMB4
Protein Name	Laminin subunit beta-4 (Laminin beta-1-related protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	LAMB4 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	193kD
Cell Pathway	Secreted, extracellular space, extracellular matrix, basement membrane.
Tissue Specificity	Placenta,Testis,
Function	domain:Domains VI and IV are globular.,domain:The alpha-helical domains I and II are thought to interact with other laminin chains to form a coiled coil structure.,function:Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components.,similarity:Contains 1 laminin IV type B domain.,similarity:Contains 1 laminin N-terminal domain.,similarity:Contains 13 laminin EGF-like domains.,subunit:Laminin is a complex glycoprotein, consisting of three different polypeptide chains (alpha, beta, gamma), which are bound to each other by disulfide bonds into a cross-shaped molecule comprising one long and three short arms with globules at each end.,
Background	domain:Domains VI and IV are globular.,domain:The alpha-helical domains I and II are thought to interact with other laminin chains to form a coiled coil structure.,function:Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during



## UpingBio technology Co.,Ltd

© Tel: 400-999-8863 ■ Email:UpingBio@163.com



embryonic development by interacting with other extracellular matrix
components., similarity: Contains 1 Iaminin IV type B domain., similarity: Contains 1
laminin N-terminal domain.,similarity:Contains 13 Iaminin EGF-like
domains., subunit: Laminin is a complex glycoprotein, consisting of three different
polypeptide chains (alpha, beta, gamma), which are bound to each other by
disulfide bonds into a cross-shaped molecule comprising one long and three shor
arms with globules at each end.,

matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images