





## **DLL1 Polyclonal Antibody**

Catalog No	YP-Ab-06840
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	DLL1 UNQ146/PRO172
Protein Name	Delta-like protein 1 (Drosophila Delta homolog 1) (Delta1) (H-Delta-1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	DLL1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Observed Barid	79kD
Cell Pathway	79kD  Apical cell membrane ; Single-pass type I membrane protein . Cell junction, adherens junction . Membrane raft . Distributed around adherens junction in the apical endfeet through interactions with MAGI1
	Apical cell membrane ; Single-pass type I membrane protein . Cell junction, adherens junction . Membrane raft . Distributed around adherens junction in the
Cell Pathway	Apical cell membrane; Single-pass type I membrane protein. Cell junction, adherens junction. Membrane raft. Distributed around adherens junction in the apical endfeet through interactions with MAGI1  Expressed in heart and pancreas, with lower expression in brain and muscle and
Cell Pathway  Tissue Specificity	Apical cell membrane; Single-pass type I membrane protein. Cell junction, adherens junction. Membrane raft. Distributed around adherens junction in the apical endfeet through interactions with MAGI1  Expressed in heart and pancreas, with lower expression in brain and muscle and almost no expression in placenta, lung, liver and kidney.  function:Acts as a ligand for Notch receptors. Blocks the differentiation of progenitor cells into the B-cell lineage while promoting the emergence of a population of cells with the characteristics of a T-cell/NK-cell precursor.,PTM:Ubiquitinated by MIB (MIB1 or MIB2), leading to its endocytosis and subsequent degradation.,similarity:Contains 1 DSL domain.,similarity:Contains 8 EGF-like domains.,subunit:Interacts with Notch receptors.,tissue specificity:Expressed in heart and pancreas, with lower expression in brain and muscle and almost no expression in placenta, lung, liver



## UpingBio technology Co.,Ltd

© Tel: 400-999-8863 ■ Email:UpingBio@163.com

Website:	www.upingBio.com	•		
Website.	www.upingbio.com			

matters	needing
attentio	

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images