



FLIP1 Polyclonal Antibody

Catalog No	YP-Ab-06848
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	FILIP1 KIAA1275
Protein Name	Filamin-A-interacting protein 1 (FILIP)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	FLIP1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	133kD
Cell Pathway	Cytoplasm, cytoskeleton .
Tissue Specificity	Moderately expressed in adult heart and brain. Weakly expressed in lung, skeletal muscle, ovary, testis, kidney, and fetal brain, and hardly detectable in liver, pancreas, spleen, and fetal liver. Within brain, moderate expression is found in amygdala and caudate nucleus.
Function	function:By acting through a filamin-A/F-actin axis, it controls the start of neocortical cell migration from the ventricular zone. May be able to induce the degradation of filamin-A.,similarity:Belongs to the FILIP1 family.,subunit:Interacts with FLNA.,tissue specificity:Moderately expressed in adult heart and brain. Weakly expressed in lung, skeletal muscle, ovary, testis, kidney, and fetal brain, and hardly detectable in liver, pancreas, spleen, and fetal liver. Within brain, moderate expression is found in amygdala and caudate nucleus.,
Background	function:By acting through a filamin-A/F-actin axis, it controls the start of neocortical cell migration from the ventricular zone. May be able to induce the degradation of filamin-A.,similarity:Belongs to the FILIP1 family.,subunit:Interacts with FLNA.,tissue specificity:Moderately expressed in adult heart and brain. Weakly expressed in lung, skeletal muscle, ovary, testis, kidney, and fetal brain, and hardly detectable in liver, pancreas, spleen, and fetal liver. Within brain,

moderate expression is found in amygdala and caudate nucleus.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images