







## FKBP4 Polyclonal Antibody

Catalog No	YP-Ab-06857
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	FKBP4 FKBP52
Protein Name	Peptidyl-prolyl cis-trans isomerase FKBP4 (PPlase FKBP4) (EC 5.2.1.8) (51 kDa FK506-binding protein) (FKBP51) (52 kDa FK506-binding protein) (52 kDa FKBP) (FKBP-52) (59 kDa immunophilin) (p59) (FK506-
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	FKBP4 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	50kD
Cell Pathway	Cytoplasm, cytosol . Mitochondrion . Nucleus . Cytoplasm, cytoskeleton . Cell projection, axon . Shuttles from mitochondria to nucleus; co-localizes in mitochondria with the glucocorticoid receptor (PubMed:21730050). Colocalized with MAPT/TAU in the distal part of the primary cortical neurons (By similarity)
Tissue Specificity	Widely expressed.
Function	catalytic activity:Peptidylproline (omega=180) = peptidylproline (omega=0).,function:Component of unactivated mammalian steroid receptor complexes that sediment at 8-10 S. May have a rotamase activity. May play a role in the intracellular trafficking of heterooligomeric forms of steroid hormone receptors.,PTM:Phosphorylation by CK2 results in loss of HSP90 binding activity (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR.,sequence caution:Wrong choice of frame.,similarity:Contains 2 PPlase FKBP-type domains.,similarity:Contains 3 TPR repeats.,subunit:Interacts with NR3C1 and dynein (By similarity). Associates with HSP90 and HSP70 in unactivated steroid hormone receptor complexes. Also interacts with peroxisomal phytanoyl-CoA alpha-hydroxylase (PHYH). Interacts with HSF1 in the HSP90 complex.,tissue specificity:Widely expressed.,



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 🛎 Email:UpingBio@163.com

Website: www.upingBio.com

Background
------------

The protein encoded by this gene is a member of the immunophilin protein family, which play a role in immunoregulation and basic cellular processes involving protein folding and trafficking. This encoded protein is a cis-trans prolyl isomerase that binds to the immunosuppressants FK506 and rapamycin. It has high structural and functional similarity to FK506-binding protein 1A (FKBP1A), but unlike FKBP1A, this protein does not have immunosuppressant activity when complexed with FK506. It interacts with interferon regulatory factor-4 and plays an important role in immunory gene expression in B and T lymphocytes. This encoded protein is known to associate with phytanoyl-CoA alpha-hydroxylase. It can also associate with two heat shock proteins (hsp90 and hsp70) and thus may play a role in the intracellular trafficking of hetero-oligomeric forms of the steroid hormone receptors. This protein corr

## matters needing attention

Avoid repeated freezing and thawing!

## **Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images