





## PARP6 Polyclonal Antibody

| Catalog No                           | YP-Ab-06878   |
|--------------------------------------|---|
| Isotype                              | IgG   |
| Reactivity                           | Human;Mouse   |
| Applications                         | WB;ELISA  |
| Gene Name                            | PARP6   |
| Protein Name                         | Poly [ADP-ribose] polymerase 6 (PARP-6) (EC 2.4.2.30) (ADP-ribosyltransferase diphtheria toxin-like 17) (ARTD17)  |
| Immunogen                            | Synthesized peptide derived from part region of human protein   |
| Specificity                          | PARP6 Polyclonal Antibody detects endogenous levels of protein.   |
| Formulation                          | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  |
| Source                               | Polyclonal, Rabbit,IgG  |
| Purification                         | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.   |
| Dilution                             | WB 1:500-2000 ELISA 1:5000-20000  |
| Concentration                        | 1 mg/ml   |
| Purity                               | ≥90%  |
| Storage Stability                    | -20°C/1 year  |
| Synonyms                             |   |
| Observed Band                        | 69kD  |
| Cell Pathway                         |   |
| Tissue Specificity                   | Coronary artery, Embryo, Kidney, PCR rescued clones, PNS, Terato  |
| Function                             | catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,similarity:Contains 1 PARP catalytic domain.,                                       |
|                                      |   |
| Background                           | catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,similarity:Contains 1 PARP catalytic domain.,                                       |
| Background matters needing attention | catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,similarity:Contains 1 PARP catalytic domain.,  Avoid repeated freezing and thawing! |
| matters needing                      |   |



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com

| Website: www.upingBio.com |  |
|---------------------------|--|

| Products Images |
|-----------------|
|                 |
|                 |
|                 |
|                 |