







PI16 Polyclonal Antibody

IgG Human;Mouse WB;ELISA PI16 CRISP9 PSPBP PSEC0164 UNQ289/PRO328 Peptidase inhibitor 16 (PI-16) (Cysteine-rich secretory protein 9) (CRISP-9) (PSP94-binding protein) Synthesized peptide derived from part region of human protein PI16 Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Human; Mouse WB; ELISA PI16 CRISP9 PSPBP PSEC0164 UNQ289/PRO328 Peptidase inhibitor 16 (PI-16) (Cysteine-rich secretory protein 9) (CRISP-9) (PSP94-binding protein) Synthesized peptide derived from part region of human protein PI16 Polyclonal Antibody detects endogenous levels of protein.
WB;ELISA PI16 CRISP9 PSPBP PSEC0164 UNQ289/PRO328 Peptidase inhibitor 16 (PI-16) (Cysteine-rich secretory protein 9) (CRISP-9) (PSP94-binding protein) Synthesized peptide derived from part region of human protein PI16 Polyclonal Antibody detects endogenous levels of protein.
PI16 CRISP9 PSPBP PSEC0164 UNQ289/PRO328 Peptidase inhibitor 16 (PI-16) (Cysteine-rich secretory protein 9) (CRISP-9) (PSP94-binding protein) Synthesized peptide derived from part region of human protein PI16 Polyclonal Antibody detects endogenous levels of protein.
Peptidase inhibitor 16 (PI-16) (Cysteine-rich secretory protein 9) (CRISP-9) (PSP94-binding protein) Synthesized peptide derived from part region of human protein PI16 Polyclonal Antibody detects endogenous levels of protein.
(PSP94-binding protein) Synthesized peptide derived from part region of human protein PI16 Polyclonal Antibody detects endogenous levels of protein.
PI16 Polyclonal Antibody detects endogenous levels of protein.
·
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
50kD
Secreted .
Expressed in prostate, testis, ovary and intestine. Concentrates in prostate cance patient's sera.
function:Putative serine protease inhibitor.,miscellaneous:May serve as a marker following prostatectomy for prostate cancer.,PTM:N-glycosylated.,similarity:Belongs to the CRISP family.,subunit:Interacts with PSP94/MSMB.,tissue specificity:Expressed in prostate, testis, ovary and intestine. Concentrates in prostate cencer patient's sera.,
function:Putative serine protease inhibitor.,miscellaneous:May serve as a marker following prostatectomy for prostate cancer.,PTM:N-glycosylated.,similarity:Belongs to the CRISP family.,subunit:Interacts with PSP94/MSMB.,tissue specificity:Expressed in prostate, testis, ovary and intestine. Concentrates in prostate cencer patient's sera.,



UpingBio technology Co.,Ltd

© Tel: 400-999-8863 ■ Email:UpingBio@163.com

(B) Wahsita	www.upingBio.com	•		
Website.	www.upingbio.com			

matters	needing
attentio	

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images