





LY96 Polyclonal Antibody

| Catalog No | YP-Ab-06926 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | LY96 ESOP1 MD2 |
| Protein Name | Lymphocyte antigen 96 (Ly-96) (ESOP-1) (Protein MD-2) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | LY96 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 17kD |
| Cell Pathway | Secreted, extracellular space . Secreted . Retained in the extracellular space at the cell surface by interaction with TLR4 (PubMed:10359581) |
| Tissue Specificity | Liver,Uterus, |
| Function | function:Cooperates with TLR4 in the innate immune response to bacterial lipopolysaccharide (LPS), and with TLR2 in the response to cell wall components from Gram-positive and Gram-negative bacteria. Enhances TLR4-dependent activation of NF-kappa-B. Cells expressing both MD2 and TLR4, but not TLR4 alone, respond to LPS.,PTM:N-glycosylated; high-mannose.,subunit:Heterogeneous homopolymer formed from homodimers; disulfide-linked. Belongs to the lipopolysaccharide (LPS) receptor, a multi-protein |
| | disulfide-linked. Belongs to the lipopolysaccharide (LPS) receptor, a multi-protein complex containing at least CD14, LY96 and TLR4. Binds to the extracellular domains of TLR2 and TLR4. Ligand binding induces interaction with TLR4 and oligomerization of the complex., |



UpingBio technology Co.,Ltd







[provided by RefSeq, Sep 2010],


| matters needing | |
|-----------------|--|
| attention | |

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4°over night