



TNKS1 Polyclonal Antibody

Catalog No	YP-Ab-07131
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	TNKS PARP5A PARPL TIN1 TINF1 TNKS1
Protein Name	Tankyrase-1 (TANK1) (EC 2.4.2.30) (ADP-ribosyltransferase diphtheria toxin-like 5) (ARTD5) (Poly [ADP-ribose] polymerase 5A) (TNKS-1) (TRF1-interacting ankyrin-related ADP-ribose polymerase) (Tankyras
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TNKS1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	145kD
Cell Pathway	Cytoplasm . Golgi apparatus membrane ; Peripheral membrane protein . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Nucleus, nuclear pore complex . Chromosome, telomere . Cytoplasm, cytoskeleton, spindle pole . Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4-vesicles (PubMed:22864114). A minor proportion is also found at nuclear pore complexes and around the pericentriolar matrix of mitotic centromeres (PubMed:10523501). During interphase, a small fraction of TNKS is found in the nucleus, associated with TERF1 (PubMed:12768206). Localizes to spindle poles at mitosis onset via interaction with NUMA1 (PubMed:12080061). .
Tissue Specificity	Ubiquitous; highest levels in testis.
Function	catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,function:May regulate vesicle trafficking and modulate the subcellular distribution of SLC2A4/GLUT4-vesicles. Has PARP activity and can modify TERF1, and thereby contribute to the regulation of telomere length.,PTM:ADP-ribosylated (-auto).,PTM:Phosphorylated on serine residues by MAPK kinases upon insulin stimulation.,similarity:Contains 1 PARP catalytic domain.,similarity:Contains 1 SAM (sterile alpha motif)



domain.,similarity:Contains 15 ANK repeats.,subcellular location:Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4-vesicles. A minor proportion is also found at nuclear pore complexes and around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS is found in the nucleus, associated with TERF1.,subunit:Oligomerizes and associates

Background

catalytic activity:NAD(+) + (ADP-D-ribosyl)(n)-acceptor = nicotinamide + (ADP-D-ribosyl)(n+1)-acceptor.,function:May regulate vesicle trafficking and modulate the subcellular distribution of SLC2A4/GLUT4-vesicles. Has PARP activity and can modify TERF1, and thereby contribute to the regulation of telomere length.,PTM:ADP-ribosylated (-auto).,PTM:Phosphorylated on serine residues by MAPK kinases upon insulin stimulation.,similarity:Contains 1 PARP catalytic domain.,similarity:Contains 1 SAM (sterile alpha motif) domain.,similarity:Contains 15 ANK repeats.,subcellular location:Associated with the Golgi and with juxtannuclear SLC2A4/GLUT4-vesicles. A minor proportion is also found at nuclear pore complexes and around the pericentriolar matrix of mitotic centromeres. During interphase, a small fraction of TNKS is found in the nucleus, associated with TERF1.,subunit:Oligomerizes and associates with TNKS2. Interacts with the cytoplasmic domain of LNPEP/Otase in SLC2A4/GLUT4-vesicles. Binds to the N-terminus of telomeric TERF1 via the ANK repeats. Found in a complex with POT1; TERF1 and TINF2.,tissue specificity:Ubiquitous; highest levels in testis.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images