



NDP Polyclonal Antibody

Catalog No	YP-Ab-07152
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	NDP EVR2
Protein Name	Norrie (Norrie disease protein) (X-linked exudative vitreoretinopathy 2 protein)
Immunogen	Synthesized peptide derived from human protein . at AA range: 40-120
Specificity	NDP Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	14kD
Cell Pathway	Secreted .
Tissue Specificity	Expressed in the outer nuclear, inner nuclear and ganglion cell layers of the retina, and in fetal and adult brain.
Function	disease:Defects in NDP are the cause of Norrie disease (ND) [MIM:310600]; also known as atrophio bulborum hereditaria or Episkopi blindness. ND is a recessive disorder characterized by very early childhood blindness due to degenerative and proliferative changes of the neuroretina. Approximately 50% of patients show some form of progressive mental disorder, often with psychotic features, and about one-third of patients develop sensorineural deafness in the second decade. In addition, some patients have more complex phenotypes, including growth failure and seizure..disease:Defects in NDP are the cause of vitreoretinopathy exudative type 2 (EVR2) [MIM:305390]. EVR2 is a disorder of the retinal vasculature characterized by an abrupt cessation of growth of peripheral capillaries, leading to an avascular peripheral retina. This may lead to compensatory retinal neovascularization, which is thou
Background	This gene encodes a secreted protein with a cystein-knot motif that activates the Wnt/beta-catenin pathway. The protein forms disulfide-linked oligomers in the extracellular matrix. Mutations in this gene result in Norrie disease and X-linked



exudative vitreoretinopathy. [provided by RefSeq, Feb 2009],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night