







OR4A5 Polyclonal Antibody

Catalog No YP-Ab-07478 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name OR4A5 Protein Name Olfactory receptor 4A5 (Olfactory receptor OR11-111) Immunogen Synthesized peptide derived from part region of human protein Specificity OR4A5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Include the Captor of Ca		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name OR4A5 Protein Name Olfactory receptor 4A5 (Olfactory receptor OR11-111) Immunogen Synthesized peptide derived from part region of human protein Specificity OR4A5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor , similarity:Belongs to the G-protein coupled receptor 1 family. Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are nembers of a large family of G-protein-coupled receptors (GPCR) varising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms. In olfactory receptor genes and proteins for this organisms is independent of other organisms. Growded by	Catalog No	YP-Ab-07478
Applications WB;ELISA Gene Name OR4A5 Protein Name Olfactory receptor 4A5 (Olfactory receptor OR11-111) Immunogen Synthesized peptide derived from part region of human protein Specificity OR4A5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens olfactory receptor receptor sinteract with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor semily is the largest in the genome. The nomenclature assigned to the olfactory receptor family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Isotype	IgG
Gene Name OR4A5 Protein Name Olfactory receptor 4A5 (Olfactory receptor OR11-111) Immunogen Synthesized peptide derived from part region of human protein Specificity OR4A5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor roteins are membres of a large family of G-protein-coupled receptors of a receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organisms is independent of other organisms, provided by	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 4A5 (Olfactory receptor OR11-111) Immunogen Synthesized peptide derived from part region of human protein Specificity OR4A5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a remembers of a large family of G-protein-coupled receptors of a responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor sene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organisms is independent of other organisms. [provided by	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein	Gene Name	OR4A5
Specificity OR4A5 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Background 0ffactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Protein Name	Olfactory receptor 4A5 (Olfactory receptor OR11-111)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Specificity	OR4A5 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor "similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors care members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of β-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Concentration	1 mg/ml
Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Synonyms	
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Background olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Observed Band	34kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family., Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background Olfactory receptor family 4 subfamily A member 5(OR4A5) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Tissue Specificity	
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by	Function	
	Background	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by



UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🛎 Email:UpingBio@163.com






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night