





## O52L2 Polyclonal Antibody

IgG Human;Rat;Mouse;  WB;ELISA  OR52L2P OR52L2  Putative olfactory receptor 52L2 (Olfactory receptor OR11-74)  Synthesized peptide derived from human protein . at AA range: 190-270  O52L2 Polyclonal Antibody detects endogenous levels of protein.  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml  ≥90%
Human;Rat;Mouse;  WB;ELISA  OR52L2P OR52L2  Putative olfactory receptor 52L2 (Olfactory receptor OR11-74)  Synthesized peptide derived from human protein . at AA range: 190-270  O52L2 Polyclonal Antibody detects endogenous levels of protein.  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
WB;ELISA  OR52L2P OR52L2  Putative olfactory receptor 52L2 (Olfactory receptor OR11-74)  Synthesized peptide derived from human protein . at AA range: 190-270  O52L2 Polyclonal Antibody detects endogenous levels of protein.  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
OR52L2P OR52L2 Putative olfactory receptor 52L2 (Olfactory receptor OR11-74) Synthesized peptide derived from human protein . at AA range: 190-270 O52L2 Polyclonal Antibody detects endogenous levels of protein. Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Polyclonal, Rabbit,IgG The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. WB 1:500-2000 ELISA 1:5000-20000
Putative olfactory receptor 52L2 (Olfactory receptor OR11-74)  Synthesized peptide derived from human protein . at AA range: 190-270  O52L2 Polyclonal Antibody detects endogenous levels of protein.  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
Synthesized peptide derived from human protein . at AA range: 190-270  O52L2 Polyclonal Antibody detects endogenous levels of protein.  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
O52L2 Polyclonal Antibody detects endogenous levels of protein.  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
Polyclonal, Rabbit,IgG  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
affinity-chromatography using epitope-specific immunogen.  WB 1:500-2000 ELISA 1:5000-20000  1 mg/ml
1 mg/ml
≥90%
-20°C/1 year
35kD
Cell membrane; Multi-pass membrane protein.
caution:Could be the product of a pseudogene.,caution:It is uncertain whether Met-1 or Met-16 is the initiator.,function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the second signal of the recognition of the largest in the second signal of the receptor gene family is the largest in the second signal of the receptor gene family is the largest in the nose, to initiate a neuronal response to the neuronal receptor second signal of the receptor second signal of the neuronal receptor second s



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com




matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night