





## SYFB Polyclonal Antibody

Catalog NoYP-Ab-07800IsotypeIgGReactivityHuman;MouseApplicationsWB;ELISAGene NameFARSB FARSLB FRSB HSPC173Protein NamePhenylalaninetRNA ligase beta subunit (EC 6.1.1.20) (Phenylalanyl-tRNA
Reactivity Human;Mouse  Applications WB;ELISA  Gene Name FARSB FARSLB FRSB HSPC173  Protein Name PhenylalaninetRNA ligase beta subunit (EC 6.1.1.20) (Phenylalanyl-tRNA
Applications WB;ELISA  Gene Name FARSB FARSLB FRSB HSPC173  Protein Name PhenylalaninetRNA ligase beta subunit (EC 6.1.1.20) (Phenylalanyl-tRNA
Gene Name FARSB FARSLB FRSB HSPC173  Protein Name PhenylalaninetRNA ligase beta subunit (EC 6.1.1.20) (Phenylalanyl-tRNA
Protein Name PhenylalaninetRNA ligase beta subunit (EC 6.1.1.20) (Phenylalanyl-tRNA
Protein Name PhenylalaninetRNA ligase beta subunit (EC 6.1.1.20) (Phenylalanyl-tRNA
synthetase beta subunit) (PheRS)
Immunogen Synthesized peptide derived from part region of human protein
Specificity SYFB Polyclonal Antibody detects endogenous levels of protein.
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source Polyclonal, Rabbit,IgG
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b> WB 1:500-2000 ELISA 1:5000-20000
Concentration 1 mg/ml
Purity ≥90%
Storage Stability -20°C/1 year
Synonyms
Observed Band 64kD
Cell Pathway Cytoplasm .
Tissue Specificity Lung, Testis, Umbilical cord blood,
Function  catalytic activity:ATP + L-phenylalanine + tRNA(Phe) = AMP + diphosphate + L-phenylalanyl-tRNA(Phe).,similarity:Belongs to the phenylalanyl-tRNA synthetase beta chain family. Type 2 subfamily.,subunit:Tetramer of two alpha and two beta chains.,
This gene encodes a highly conserved enzyme that belongs to the aminoacyl-tRNA synthetase class IIc subfamily. This enzyme comprises the regulatory beta subunits that form a tetramer with two catalytic alpha subunits. Ithe presence of ATP, this tetramer is responsible for attaching L-phenylalanine the terminal adenosine of the appropriate tRNA. A pseudogene located on chromosome 10 has been identified. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jan 2015],
matters needing Avoid repeated freezing and thawing! attention



## UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images