

Website: www.upingBio.com

SUH Polyclonal Antibody

Catalog No	YP-Ab-07839
Isotype	lgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	RBPJ IGKJRB IGKJRB1 RBPJK RBPSUH
Protein Name	Recombining binding protein suppressor of hairless (CBF-1) (J kappa-recombination signal-binding protein) (RBP-J kappa) (RBP-J) (RBP-JK) (Renal carcinoma antigen NY-REN-30)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SUH Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	55kD
Cell Pathway	Nucleus. Cytoplasm. Mainly nuclear, upon interaction with RITA/C12orf52, translocates to the cytoplasm, down-regulating the Notch signaling pathway.
Tissue Specificity	Eye,Placenta,Renal cell carcinoma,T
Function	caution:Despite some similarity with the "phage" integrase family, it has no recombinase activity.,function:Transcriptional regulator that plays a central role in Notch signaling, a signaling pathway involved in cell-cell communication that regulates a broad spectrum of cell-fate. determinations. Acts as a transcriptional repressor when it is not associated with Notch proteins. When associated with some Notch protein, it acts as a transcriptional activator that activates transcription of Notch target genes. Probably represses or activates transcription via the recruitment of chromatin remodeling complexes containing histone deacetylase or histone acetylase proteins, respectively. Specifically binds to the immunoglobulin kappa-type J segment recombination signal sequence.,similarity:Belongs to the Su(H) family.,similarity:Contains 1 IPT/TIG domain.,subunit:Interacts with activated NOTCH1,
Background	The protein encoded by this gene is a transcriptional regulator important in the Notch signaling pathway. The encoded protein acts as a repressor when not



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 💌 Email:UpingBio@163.com

Website: www.upingBio.com

bound to Notch proteins and an activator when bound to Notch proteins. It is
thought to function by recruiting chromatin remodeling complexes containing
histone deacetylase or histone acetylase proteins to Notch signaling pathway
genes. Several transcript variants encoding different isoforms have been found for
this gene, and several pseudogenes of this gene exist on chromosome 9.
[provided by RefSeq, Oct 2013],matters needing
attentionAvoid repeated freezing and thawing!Usage suggestionsThis product can be used in immunological reaction related experiments. For
more information, please consult technical personnel.

Products Images