





MMP21 rabbit pAb

| Catalog No | YP-Ab-08055 |
|--------------------|--|
| Isotype | IgG |
| Reactivity | Human; Mouse |
| Applications | WB |
| Gene Name | MMP21 |
| Protein Name | MMP21 |
| Immunogen | Synthesized peptide derived from human MMP21 AA range: 425-475 |
| Specificity | This antibody detects endogenous levels of MMP21 at Human/Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1:500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) |
| Observed Band | 60kD |
| Cell Pathway | Secreted . |
| Tissue Specificity | Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. |
| Function | cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines., |
| Background | This gene encodes a member of the matrix metalloproteinase family. Proteins in this family are involved in the breakdown of extracellular matrix for both normal physiological processes, such as embryonic development, reproduction, and |



UpingBio technology Co.,Ltd

C Tel: 400-999-8863 ■ Email:UpingBio@163.com



tissue remodeling, and disease processes, such as asthma and tumor metastasis. The encoded protein may play an important role in embryogenesis, particularly in neuronal cells, as well as in lymphocyte development and survival. [provided by RefSeq, May 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night