






TRI50 rabbit pAb

Catalog No	YP-Ab-08254
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TRIM50 TRIM50A
Protein Name	TRI50
Immunogen	Synthesized peptide derived from human TRI50 AA range: 312-362
Specificity	This antibody detects endogenous levels of TRI50 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year Cytoplasm . Localizes mainly into discrete cytoplasmic punctuate structures
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Cytoplasm . Localizes mainly into discrete cytoplasmic punctuate structures heterogeneous in size and shape containing polyubiquitinated proteins function:E3 ubiquitin-protein ligase.,PTM:Auto-ubiquitinated.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Cytoplasm . Localizes mainly into discrete cytoplasmic punctuate structures heterogeneous in size and shape containing polyubiquitinated proteins function:E3 ubiquitin-protein ligase.,PTM:Auto-ubiquitinated.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,subunit:Can form dimers and trimers. Interacts with several E2 ubiquitin-conjugating enzymes, including UBE2L6, UBE2E1, UBE2E3. No
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Cytoplasm . Localizes mainly into discrete cytoplasmic punctuate structures heterogeneous in size and shape containing polyubiquitinated proteins function:E3 ubiquitin-protein ligase.,PTM:Auto-ubiquitinated.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,subunit:Can form dimers and trimers. Interacts with several E2 ubiquitin-conjugating enzymes, including UBE2L6, UBE2E1, UBE2E3. No
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Cytoplasm . Localizes mainly into discrete cytoplasmic punctuate structures heterogeneous in size and shape containing polyubiquitinated proteins function:E3 ubiquitin-protein ligase.,PTM:Auto-ubiquitinated.,similarity:Belongs to the TRIM/RBCC family.,similarity:Contains 1 B box-type zinc finger.,similarity:Contains 1 B30.2/SPRY domain.,similarity:Contains 1 RING-type zinc finger.,subunit:Can form dimers and trimers. Interacts with several E2 ubiquitin-conjugating enzymes, including UBE2L6, UBE2E1, UBE2E3. No interaction with UBE2H.,





Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night