


K1456 rabbit pAb

| Catalog No | YP-Ab-08284 |
|---|---|
| Isotype | IgG |
| Reactivity | Human; Mouse |
| Applications | WB |
| Gene Name | KIAA1456 C8orf79 |
| Protein Name | K1456 |
| Immunogen | Synthesized peptide derived from human K1456 AA range: 140-190 |
| Specificity | This antibody detects endogenous levels of K1456 at Human/Mouse |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1: 500-2000 |
| Concentration | 1 mg/ml |
| Concentration | · · · · g · · · · |
| Purity | ≥90% |
| | |
| Purity | ≥90% |
| Purity Storage Stability | ≥90% |
| Purity Storage Stability Synonyms | ≥90% |
| Purity Storage Stability Synonyms Observed Band | ≥90% -20°C/1 year |
| Purity Storage Stability Synonyms Observed Band Cell Pathway | ≥90% -20°C/1 year nucleus,cytoplasm, |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity | ≥90% -20°C/1 year nucleus,cytoplasm, Down-regulated in breast, bladder, colorectal, cervix and testicular carcinomas. |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | ≥90% -20°C/1 year nucleus,cytoplasm, Down-regulated in breast, bladder, colorectal, cervix and testicular carcinomas. |
| Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing | ≥90% -20°C/1 year nucleus,cytoplasm, Down-regulated in breast, bladder, colorectal, cervix and testicular carcinomas. similarity:Belongs to the methyltransferase superfamily., |



Products Images





Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night