


S22AN rabbit pAb

Catalog No	YP-Ab-08291
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	SLC22A23 C6orf85
Protein Name	S22AN
Immunogen	Synthesized peptide derived from human S22AN AA range: 600-650
Specificity	This antibody detects endogenous levels of S22AN at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
	· ··· 9 ·····
Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Membrane ; Multi-pass membrane protein . similarity:Belongs to the major facilitator superfamily. Organic cation transporter
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Membrane; Multi-pass membrane protein. similarity:Belongs to the major facilitator superfamily. Organic cation transporter family., SLC22A23 belongs to a large family of transmembrane proteins that function as uniporters, symporters, and antiporters to transport organic ions across cell membranes (Jacobsson et al., 2007 [PubMed 17714910]).[supplied by OMIM,
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Membrane ; Multi-pass membrane protein . similarity:Belongs to the major facilitator superfamily. Organic cation transporter family., SLC22A23 belongs to a large family of transmembrane proteins that function as uniporters, symporters, and antiporters to transport organic ions across cell membranes (Jacobsson et al., 2007 [PubMed 17714910]).[supplied by OMIM, Mar 2008],



Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night