





YN3572

## IKZF5 rabbit pAb

Catalog No	YP-Ab-08414
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	IKZF5 ZNFN1A5
Protein Name	IKZF5
Immunogen	Synthesized peptide derived from human IKZF5 AA range: 210-260
Specificity	This antibody detects endogenous levels of IKZF5 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in brain, heart, skeletal muscle, kidney, and liver. Expressed in the hematopoietic cell lines MOLT-4, NALM-6 and K-562. Highly expressed in THP-1 and M-07e cell lines, which have characteristics of myeloid and early megakaryocytic cells respectively.
Function	domain:C-terminal zinc fingers mediate homodimerization.,domain:The N-terminal zinc fingers are involved in sequence-specific DNA binding and heterotypic associations with other family members.,function:DNA-binding protein that binds to the 5'GNNTGTNG-3' core sequence. Transcriptional repressor.,miscellaneous:'Pegasus' was the winged horse in Greek mythology.,similarity:Belongs to the Ikaros C2H2-type zinc-finger protein family.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Self-associates and interacts with other family members; IKZF1, IKZF2, IKZF3 AND IKZF4.,tissue specificity:Expressed in brain, heart, skeletal muscle, kidney, and liver. Expressed




## UpingBio technology Co.,Ltd





	in the hematopoietic cell lines MOLT-4, NALM-6 and K562. Highly expressed in THP-1 and Mo7e-IL3 cell lines, which have characteristics of myeloid and early megakaryocytic cells respectively.,
Background	Members of the Ikaros (ZNFN1A1; MIM 603023) family of transcription factors, which includes Pegasus, are expressed in lymphocytes and are implicated in the control of lymphoid development.[supplied by OMIM, Jul 2002],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4°over night