







ZUFSP rabbit pAb


Catalog No	YP-Ab-08454
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	ZUFSP C6orf113
Protein Name	ZUFSP
Immunogen	Synthesized peptide derived from human ZUFSP AA range: 89-139
Specificity	This antibody detects endogenous levels of ZUFSP at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration	1 mg/ml
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cytoplasm . Nucleus . Mostly present in the nuclear fraction. Localizes to DNA
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm . Nucleus . Mostly present in the nuclear fraction. Localizes to DNA
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm . Nucleus . Mostly present in the nuclear fraction. Localizes to DNA lesions caution:Although related to the Ufm1-specific proteases, it lacks the conserved His active site, suggesting that it is probably inactive.,similarity:Contains 3
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm . Nucleus . Mostly present in the nuclear fraction. Localizes to DNA lesions caution:Although related to the Ufm1-specific proteases, it lacks the conserved His active site, suggesting that it is probably inactive.,similarity:Contains 3
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cytoplasm . Nucleus . Mostly present in the nuclear fraction. Localizes to DNA lesions caution:Although related to the Ufm1-specific proteases, it lacks the conserved His active site, suggesting that it is probably inactive.,similarity:Contains 3 C2H2-type zinc fingers.,







Products Images



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night