







MORN2 rabbit pAb

Catalog No	YP-Ab-08589
Isotype	IgG
Reactivity	Human; Mouse
Applications	IHC;IF
Gene Name	MORN2 MOPT
Protein Name	MORN2
Immunogen	Synthesized peptide derived from human MORN2 AA range: 25-75
Specificity	This antibody detects endogenous levels of MORN2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	5000/
· arity	≥90%
Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis similarity:Contains 2 MORN repeats.,




UpingBio technology Co.,Ltd







Products Images



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).