





## CHIC2 rabbit pAb

Catalog No	YP-Ab-08647
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	CHIC2 BTL
Protein Name	CHIC2
Immunogen	Synthesized peptide derived from human CHIC2 AA range: 111-161
Specificity	This antibody detects endogenous levels of CHIC2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
	· ···· <del>g</del> ·····
Purity	≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year  Cell membrane . Cytoplasmic vesicle . Also present at a Golgi-like vesicular
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year  Cell membrane . Cytoplasmic vesicle . Also present at a Golgi-like vesicular
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90%  -20°C/1 year  Cell membrane . Cytoplasmic vesicle . Also present at a Golgi-like vesicular compartment and at scattered vesicles.  disease:A chromosomal aberration involving CHIC2 is found in a form of acute myeloid leukemia (AML). Translocation t(4;12)(q12;p13) with ETV6.,PTM:Palmitoylation in the CHIC motif is required for membrane association.,similarity:Belongs to the CHIC family.,subcellular location:Also
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90%  -20°C/1 year  Cell membrane . Cytoplasmic vesicle . Also present at a Golgi-like vesicular compartment and at scattered vesicles.  disease:A chromosomal aberration involving CHIC2 is found in a form of acute myeloid leukemia (AML). Translocation t(4;12)(q12;p13) with ETV6.,PTM:Palmitoylation in the CHIC motif is required for membrane association.,similarity:Belongs to the CHIC family.,subcellular location:Also present at a Golgi-like vesicular compartment and at scattered vesicles.,  This gene encodes a member of the CHIC family of proteins. The encoded protein contains a cysteine-rich hydrophobic (CHIC) motif, and is localized to vesicular structures and the plasma membrane. This gene is associated with some cases of



## UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night