







BAHD1 rabbit pAb


Catalog No	YP-Ab-08718
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	BAHD1 KIAA0945
Protein Name	BAHD1
Immunogen	Synthesized peptide derived from human BAHD1 AA range: 214-264
Specificity	This antibody detects endogenous levels of BAHD1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Nucleus . Chromosome . Localizes to heterochromatin and inactive X
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Nucleus . Chromosome . Localizes to heterochromatin and inactive X
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Nucleus . Chromosome . Localizes to heterochromatin and inactive X chromosome. Colocalizes with histone H3 trimethylated at 'Lys-27' (H3K27me3).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Nucleus . Chromosome . Localizes to heterochromatin and inactive X chromosome. Colocalizes with histone H3 trimethylated at 'Lys-27' (H3K27me3).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Nucleus . Chromosome . Localizes to heterochromatin and inactive X chromosome. Colocalizes with histone H3 trimethylated at 'Lys-27' (H3K27me3). similarity:Contains 1 BAH domain.,







Products Images



Western blot analysis of lysates from VEC cells, primary antibody was diluted at 1:1000, 4° over night