




NAA60 rabbit pAb

Catalog No	YP-Ab-08810
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	NAA60 HAT4 NAT15 UNQ2771/PRO7155
Protein Name	NAA60
Immunogen	Synthesized peptide derived from human NAA60 AA range: 5-55
Specificity	This antibody detects endogenous levels of NAA60 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Golgi apparatus membrane ; Peripheral membrane protein ; Cytoplasmic side . Probably forms an intramembrane hairpin-like structure in the membrane. .
Tissue Specificity	
Function	function: Probable N-acetyltransferase., similarity: Belongs to the acetyltransferase family., similarity: Contains 1 N-acetyltransferase domain.,
Background	This gene encodes an enzyme that localizes to the Golgi apparatus, where it transfers an acetyl group to the N-terminus of free proteins. This enzyme acts on histones, and its activity is important for chromatin assembly and chromosome integrity. Alternative splicing and the use of alternative promoters results in multiple transcript variants. The upstream promoter is located in a differentially methylated region (DMR) and undergoes imprinting; transcript variants originating from this position are expressed from the maternal allele. [provided by RefSeq, Nov 2015],
matters needing attention	Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western blot analysis of lysates from HeLa cells,
primary antibody was diluted at 1:1000, 4° over night