







UBFL1 rabbit pAb


Catalog No	YP-Ab-08869
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	UBTFL1 C11orf27
Protein Name	UBFL1
Immunogen	Synthesized peptide derived from human UBFL1 AA range: 223-273
Specificity	This antibody detects endogenous levels of UBFL1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cytoplasm . Nucleus . Mainly cytoplasmic from the 4-cell stage to the morula
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm . Nucleus . Mainly cytoplasmic from the 4-cell stage to the morula
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm . Nucleus . Mainly cytoplasmic from the 4-cell stage to the morula
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm . Nucleus . Mainly cytoplasmic from the 4-cell stage to the morula stage. Becomes nuclear at the blastocyst stage UBTFL1 is a preimplantation-specific gene and is involved in early development, implantation, and embryonic stem (ES) cell derivation (summary by Yamada et
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cytoplasm . Nucleus . Mainly cytoplasmic from the 4-cell stage to the morula stage. Becomes nuclear at the blastocyst stage UBTFL1 is a preimplantation-specific gene and is involved in early development, implantation, and embryonic stem (ES) cell derivation (summary by Yamada et al., 2010 [PubMed 19915186]).[supplied by OMIM, Jan 2011],







Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night