




NP1L1 rabbit pAb

Catalog No	YP-Ab-08919
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	NAP1L1 NRP
Protein Name	NP1L1
Immunogen	Synthesized peptide derived from human NP1L1 AA range: 272-322
Specificity	This antibody detects endogenous levels of NP1L1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus . Melanosome . Cytoplasm . Identified by mass spectrometry in melanosome fractions from stage I to stage IV.
Tissue Specificity	Ubiquitously expressed.
Function	domain: The acidic domains are probably involved in the interaction with histones., function: May be involved in modulating chromatin formation and contribute to regulation of cell proliferation., similarity: Belongs to the nucleosome assembly protein (NAP) family., subcellular location: Identified by mass spectrometry in melanosome fractions from stage I to stage IV., tissue specificity: Ubiquitously expressed.,
Background	This gene encodes a member of the nucleosome assembly protein (NAP) family. This protein participates in DNA replication and may play a role in modulating chromatin formation and contribute to the regulation of cell proliferation. Alternative splicing results in multiple transcript variants encoding different isoforms; however, not all have been fully described. [provided by RefSeq, Apr 2015],
matters needing attention	Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night