


TFPT rabbit pAb

| Catalog No | YP-Ab-08959 |
|---------------------------|--|
| Isotype | IgG |
| Reactivity | Human; Mouse;Rat |
| Applications | WB |
| Gene Name | TFPT INO80F |
| Protein Name | TFPT |
| Immunogen | Synthesized peptide derived from human TFPT AA range: 47-97 |
| Specificity | This antibody detects endogenous levels of TFPT at Human/Mouse/Rat |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. |
| Dilution | WB 1: 500-2000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | |
| Cell Pathway | Nucleus . |
| Tissue Specificity | |
| Function | disease:A chromosomal aberration involving TFPT is a cause of pre-B-cell acute lymphoblastic leukemia (B-ALL). Inversion inv(19)(p13;q13) with TCF3.,subunit:Interacts with NOL3 (By similarity). Component of the chromatin-remodeling INO80 complex, at least composed of ACTL6A, ACTR5, ACTR8, RVBL1, RVBL2, INO80, INO80B, INO80C, INO80D and INO80E., |
| Background | |
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |



Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night