

KILIN rabbit pAb

Catalog No	YP-Ab-09049
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	KLLN
Protein Name	KILIN
Immunogen	Synthesized peptide derived from human KILIN AA range: 92-142
Specificity	This antibody detects endogenous levels of KILIN at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
•	
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year Nucleus .
Synonyms Observed Band	
Synonyms Observed Band Cell Pathway	
Synonyms Observed Band Cell Pathway Tissue Specificity	Nucleus . function:DNA-binding protein involved in S phase checkpoint control-coupled apoptosis by mediating p53/TP53-induced apoptosis. Has the ability to inhibit DNA synthesis and S phase arrest coupled to apoptosis. Has affinity to both double- and single-stranded DNA.,induction:by p53/TP53; direct transcription
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Nucleus . function:DNA-binding protein involved in S phase checkpoint control-coupled apoptosis by mediating p53/TP53-induced apoptosis. Has the ability to inhibit DNA synthesis and S phase arrest coupled to apoptosis. Has affinity to both double- and single-stranded DNA.,induction:by p53/TP53; direct transcription target of p53/TP53., The protein encoded by this intronless gene is found in the nucleus, where it can inhibit DNA synthesis and promote S phase arrest coupled to apoptosis. The expression of this DNA binding protein is upregulated by transcription factor p53.



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Jarkat cells, primary antibody was diluted at 1:1000, 4° over night