

FAF2 rabbit pAb

YP-Ab-09055
IgG
Human; Mouse;Rat
WB
FAF2 ETEA KIAA0887 UBXD8 UBXN3B
FAF2
Synthesized peptide derived from human FAF2 AA range: 120-170
This antibody detects endogenous levels of FAF2 at Human/Mouse/Rat
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
WB 1: 500-2000
1 mg/ml
≥90%
-20°C/1 year
Cytoplasm . Lipid droplet . Endoplasmic reticulum .
Broadly expressed, with highest levels in brain.
caution:It seems odd to name this protein FAS-associated factor, but not have any mention of this in the CC lines.,miscellaneous:Up-regulated in T-cells and eosinophils from patients with atopic dermatitis.,similarity:Contains 1 UBA domain.,similarity:Contains 1 UBX domain.,tissue specificity:Broadly expressed, with highest levels in brain.,
The protein encoded by this gene is highly expressed in peripheral blood of patients with atopic dermatitis (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients. [provided by RefSeq, Jul 2008],
Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night