





PHF11 rabbit pAb

Catalog No YP-Ab-09109 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name PHF11 BCAP Protein Name PHF11 Immunogen Synthesized peptide derived from human PHF11 AA range: 232-282 Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal itssues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products.A number of isoforms may be produced, polymorphism-Variation in PHF11 seems to be associat		
Reactivity Human; Mouse;Rat Applications WB Gene Name PHF11 BCAP Protein Name PHF11 Immunogen Synthesized peptide derived from human PHF11 AA range: 232-282 Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus. Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthmas, similarity. Contains 1 PHD-type zinc finger. Submit. Interacts with BRCA1. Lissues specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Background This gene has been identified in some studies as a candidate gene for asthmas. Naturally-occurring erealthought transcriptor may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants, [provided by RefSeq, Mar 2016],	Catalog No	YP-Ab-09109
Applications WB Gene Name PHF11 BCAP Protein Name PHF11 Immunogen Synthesized peptide derived from human PHF11 AA range: 232-282 Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells Expressed at low levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal itssues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism's variation in PHF11 seems to be associated with propensity to atopy and asthma. similarity Contains 1 PHD-type zinc finger. Submit. Interacts with BRCA1 tissues pested (itssues hese in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcriptor may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016].	Isotype	IgG
Gene Name PHF11 BCAP Protein Name PHF11 Immunogen Synthesized peptide derived from human PHF11 AA range: 232-282 Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells: Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: An umber of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma. similarity: Contains 1 PHD-type zinc finger, subunit. Interacts with BTCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Plackground This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene nace been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],	Reactivity	Human; Mouse;Rat
Protein Name PHF11 Immunogen Synthesized peptide derived from human PHF11 AA range: 232-282 Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: An umber of isoforms may be produced polymorphism Variation in PHF11 seems to be associated with propensity to atopy and asthma. similarity:Contains 1 PHD-type zinc finger, subunit:Interacts with BRCA1. tissue specificity:Expressed in all normal issues tested, including lung, testis, small intestine, breast, liver and placenta. Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], Moritary Phone P	Applications	WB
Immunogen Synthesized peptide derived from human PHF11 AA range: 232-282 Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma, similarity. Contains 1 PHD-type zinc finger, subunit: Interacts with BRCA1, tissue specificity: Expressed in all normal tissues tested, including lung a PHD (plant homeodomain) type zinc finger. This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],	Gene Name	PHF11 BCAP
Specificity This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus. Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all anormal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism/Variation in PHF11 seems to be associated with propensity to atopy and astfmar, similarity. Contains 1 PHD-type zinc linger., subunit: Interacts with BRCA1, tissue specificity. Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for astfman. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing	Protein Name	PHF11
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus. Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma. similarity: Contains 1 PHD-type zinc finger, subunit: Interacts with BRCA1, ilssue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thaving!	Immunogen	Synthesized peptide derived from human PHF11 AA range: 232-282
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma, similarity: Contains 1 PHD-type zinc inger, subunit: Interacts with BRCA1, tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],	Specificity	This antibody detects endogenous levels of PHF11 at Human/Mouse/Rat
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus. Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative producets: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma, similarity: Contains 1 PHD-type zinc finger. subunit: Interacts with BRCA1, tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma. similarity: Contains 1 PHD-type zinc finger, subunit: Interacts with BRCA1. tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Background This gene encodes a protein containing a PHD (pant) homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma, similarity: Contains 1 PHD-type zinc finger., subunit: Interacts with BRCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], Matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma., similarity: Contains 1 PHD-type zinc finger., subunit: Interacts with BRCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma. similarity: Contains 1 PHD-type zinc finger. subunit: Interacts with BRCA1. tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Observed Band Cell Pathway Nucleus Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma., similarity: Contains 1 PHD-type zinc finger., subunit: Interacts with BRCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Cell Pathway Nucleus . Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma., similarity: Contains 1 PHD-type zinc finger., subunit: Interacts with BRCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma., similarity: Contains 1 PHD-type zinc finger., subunit: Interacts with BRCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., Background This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Highly expressed in T and B-cells, as well as natural killer and mature dendritic cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products:A number of isoforms may be produced,polymorphism:Variation in PHF11 seems to be associated with propensity to atopy and asthma.,similarity:Contains 1 PHD-type zinc finger.,subunit:Interacts with BRCA1.,tissue specificity:Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Observed Band	
cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta. Function alternative products: A number of isoforms may be produced, polymorphism: Variation in PHF11 seems to be associated with propensity to atopy and asthma., similarity: Contains 1 PHD-type zinc finger., subunit: Interacts with BRCA1., tissue specificity: Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Cell Pathway	Nucleus .
produced,polymorphism:Variation in PHF11 seems to be associated with propensity to atopy and asthma.,similarity:Contains 1 PHD-type zinc finger.,subunit:Interacts with BRCA1.,tissue specificity:Expressed in all normal tissues tested, including lung, testis, small intestine, breast, liver and placenta., This gene encodes a protein containing a PHD (plant homeodomain) type zinc finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], Matters needing Avoid repeated freezing and thawing!	Tissue Specificity	cells. Expressed at higher levels in Th1 as compared to Th2 cells. Expressed at low levels in all normal tissues tested, including lung, testis, small intestine,
finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016], matters needing Avoid repeated freezing and thawing!	Function	produced,polymorphism:Variation in PHF11 seems to be associated with propensity to atopy and asthma.,similarity:Contains 1 PHD-type zinc finger.,subunit:Interacts with BRCA1.,tissue specificity:Expressed in all normal
	Background	finger. This gene has been identified in some studies as a candidate gene for asthma. Naturally-occurring readthrough transcription may occur from the upstream SETDB2 (SET domain bifurcated 2) gene to this locus. Alternative
attention		Avoid repeated freezing and thawing!



UpingBio technology Co.,Ltd


C Tel: 400-999-8863 ■ Email:UpingBio@163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from SH-SY5Y cells, primary antibody was diluted at 1:1000, 4° over night