






## TDIF1 rabbit pAb

Catalog No	YP-Ab-09172
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	DNTTIP1 C20orf167 TDIF1
Protein Name	TDIF1
Immunogen	Synthesized peptide derived from human TDIF1 AA range: 208-258
Specificity	This antibody detects endogenous levels of TDIF1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
	>000/
Purity	≥90%
Purity Storage Stability	-20°C/1 year
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  Nucleus .  function:Shown to enhance TdT activity, in vitro.,subunit:Monomer, homodimer
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  Nucleus .  function:Shown to enhance TdT activity, in vitro.,subunit:Monomer, homodimer and homotrimer. Interacts with terminal deoxynucleotidyltransferase (TdT).,  DNTTIP1 binds DNA and enhances the activity of terminal deoxynucleotidyltransferase (TDT, or DNTT; MIM 187410), a DNA polymerase that catalyzes the polymerization of DNA in the absence of a DNA template
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year  Nucleus .  function:Shown to enhance TdT activity, in vitro.,subunit:Monomer, homodimer and homotrimer. Interacts with terminal deoxynucleotidyltransferase (TdT).,  DNTTIP1 binds DNA and enhances the activity of terminal deoxynucleotidyltransferase (TDT, or DNTT; MIM 187410), a DNA polymerase that catalyzes the polymerization of DNA in the absence of a DNA template (Yamashita et al., 2001 [PubMed 11473582]).[supplied by OMIM, Mar 2008],



## **Products Images**



Western blot analysis of lysates from HEK293 cells, primary antibody was diluted at 1:1000, 4° over night