

## PSB7 rabbit pAb

Catalog No	YP-Ab-09238
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	PSMB7 Z
Protein Name	Proteasome subunit beta type-7 (EC 3.4.25.1) (Macropain chain Z) (Multicatalytic endopeptidase complex chain Z) (Proteasome subunit Z)
Immunogen	Synthesized peptide derived from human PSB7 AA range: 191-241
Specificity	This antibody detects endogenous levels of human PSB7
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
•	
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year  Cytoplasm . Nucleus . Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  Cytoplasm . Nucleus . Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9  Expressed at a low level in colonic mucosa. Up-regulated in colorectal cancer
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  Cytoplasm . Nucleus . Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9  Expressed at a low level in colonic mucosa. Up-regulated in colorectal cancer
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  Cytoplasm . Nucleus . Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9  Expressed at a low level in colonic mucosa. Up-regulated in colorectal cancer
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	-20°C/1 year  Cytoplasm . Nucleus . Translocated from the cytoplasm into the nucleus following interaction with AKIRIN2, which bridges the proteasome with the nuclear import receptor IPO9  Expressed at a low level in colonic mucosa. Up-regulated in colorectal cancer tissues.



## UpingBio technology Co.,Ltd

🕻 Tel: 400-999-8863 📾 Email:UpingBio@163.com

Website: www.upingBio.com
---------------------------

Products Images