







HUNIN rabbit pAb

Catalog No YP-Ab-09285 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name MT-RNR2 Protein Name Putative humanin peptide Immunogen Synthesized peptide derived from human HUNIN AA range: 1-24 Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band		
Reactivity Human;Rat;Mouse; Applications WB Gene Name MT-RNR2 Protein Name Putative humanin peptide Immunogen Synthesized peptide derived from human HUNIN AA range: 1-24 Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Catalog No	YP-Ab-09285
Applications WB Gene Name MT-RNR2 Protein Name Putative humanin peptide Immunogen Synthesized peptide derived from human HUNIN AA range: 1-24 Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity \$90% Storage Stability -20°C/1 year Synonyms	Isotype	IgG
Gene Name MT-RNR2 Protein Name Putative humanin peptide Immunogen Synthesized peptide derived from human HUNIN AA range: 1-24 Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Reactivity	Human;Rat;Mouse;
Protein Name Putative humanin peptide Immunogen Synthesized peptide derived from human HUNIN AA range: 1-24 Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Applications	WB
Immunogen Synthesized peptide derived from human HUNIN AA range: 1-24 Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Gene Name	MT-RNR2
Specificity This antibody detects endogenous levels of human HUNIN Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Protein Name	Putative humanin peptide
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Immunogen	Synthesized peptide derived from human HUNIN AA range: 1-24
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Specificity	This antibody detects endogenous levels of human HUNIN
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Dilution WB 1:1000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms	Dilution	WB 1:1000-2000
Storage Stability -20°C/1 year Synonyms	Concentration	1 mg/ml
Synonyms	Purity	≥90%
	Storage Stability	-20°C/1 year
Observed Band	Synonyms	
	Observed Band	

Cell Pathway Secreted . Cytoplasm . Cell projection, cilium, flagellum . Nucleus . Mitochondrion . Localizes to the sperm flagellum where it is highly concentrated in the midpiece (PubMed:20542501, PubMed:30920769). Detected in the cytoplasm and nucleus of spermatocytes and spermatids (PubMed:20542501). Also detected in sperm mitochondria (PubMed:20542501). In retinal pigment epithelium cells, detected in cytoplasm and mitochondria (PubMed:26990160).

Expressed in testis, seminal plasma and sperm (at protein level) (PubMed:20542501, PubMed:30920769). Higher seminal plasma levels are **Tissue Specificity**

associated with normospermia than with oligospermia, asthenospermia or oligoasthenospermia (at protein level) (PubMed:30920769). Higher sperm levels are associated with normospermia than with asthenospermia (at protein level) (PubMed:30920769). Expressed in retinal epithelial cells (at protein level) (PubMed:26990160). Expressed in the heart, skeletal muscle, kidney and liver. Lesser but significant expression is observed in the brain and the gastrointestinal tract. Expressed in the AD brain, where it is found in some of the large intact neurons of the occipital lobes and small and round reactive glial cells in the hippocampus.

Function



UpingBio technology Co.,Ltd

€ Tel: 400-999-8863 🛎 Email:UpingBio@163.com



Background

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images