



# F110C rabbit pAb

<b>Catalog No</b>	YP-Ab-09354
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	FAM110C
<b>Protein Name</b>	Protein FAM110C
<b>Immunogen</b>	Synthesized peptide derived from human F110C AA range: 109-159
<b>Specificity</b>	This antibody detects endogenous levels of human F110C
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . Cytoplasm, cytoskeleton, spindle pole . Nucleus . Colocalizes with microtubules during interphase (PubMed:17499476). Detected at the mitotic spindle poles (PubMed:17499476). Colocalizes with AKT1 at the cell cortex (PubMed:19698782). .
<b>Tissue Specificity</b>	Detected in stomach, thyroid, trachea, adrenal gland and testis, and at low levels in prostate, ovary, intestine, colon, spinal cord and lymph node.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images