

Tel: 400-999-8863
■ Emall:Upingbio.163.com





PTGDS rabbit pAb

Catalog No	YP-Ab-09411	
Isotype	IgG	
Reactivity	Human;Mouse;Rat	
Applications	WB	
Gene Name	PTGDS PDS	
Protein Name	Prostaglandin-H2 D-isomerase (EC 5.3.99.2) (Beta-trace protein) (Cerebrin-28) (Glutathione-independent PGD synthase) (Lipocalin-type prostaglandin-D synthase) (Prostaglandin-D2 synthase) (PGD2 synthas	
Immunogen	Synthesized peptide derived from human PTGDS AA range: 115-165	
Specificity	This antibody detects endogenous levels of human PTGDS	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.	
Dilution	WB 1:1000-2000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Observed Band		
Cell Pathway	Rough endoplasmic reticulum . Nucleus membrane . Golgi apparatus . Cytoplasm, perinuclear region . Secreted . Detected on rough endoplasmic reticulum of arachnoid and menigioma cells. Localized to the nuclear envelope, Golgi apparatus, secretory vesicles and spherical cytoplasmic structures in arachnoid trabecular cells, and to circular cytoplasmic structures in meningeal macrophages and perivascular microglial cells. In oligodendrocytes, localized to the rough endoplasmic reticulum and nuclear envelope. In retinal pigment epithelial cells, localized to distinct cytoplasmic domains including the perinuclear region. Also secreted.	
Tissue Specificity	Abundant in the brain and CNS, where it is expressed in tissues of the blood-brain barrier and secreted into the cerebro-spinal fluid. Abundantly expressed in the heart. In the male reproductive system, it is expressed in the testis, epididymis and prostate, and is secreted into the seminal fluid. Expressed in the eye and secreted into the aqueous humor. Lower levels detected in various tissue fluids such as serum, normal urine, ascitic fluid and tear fluid. Also found in a number of other organs including ovary, fimbriae of the fallopian tubes, kidney, leukocytes.	

Function

other organs including ovary, fimbriae of the fallopian tubes, kidney, leukocytes.



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Background

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images