



# SPG20 rabbit pAb

<b>Catalog No</b>	YP-Ab-09717
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	SPG20 KIAA0610 TAHCCP1
<b>Protein Name</b>	Spartin (Spastic paraplegia 20 protein) (Trans-activated by hepatitis C virus core protein 1)
<b>Immunogen</b>	Synthesized peptide derived from human SPG20 AA range: 303-353
<b>Specificity</b>	This antibody detects endogenous levels of human SPG20
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Midbody . Transiently associated with endosomes (PubMed:19580544). Colocalized with IST1 to the ends of Flemming bodies during cytokinesis (PubMed:20719964). .
<b>Tissue Specificity</b>	Ubiquitously expressed, with highest levels of expression detected in adipose tissue.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images