





MCA3 rabbit pAb

Catalog No	YP-Ab-09770	
Isotype	IgG	
Reactivity	Human;Mouse	
Applications	WB	
Gene Name	EEF1E1 AIMP3 P18	
Protein Name	Eukaryotic translation elongation factor 1 epsilon-1 (Aminoacyl tRNA synthetase complex-interacting multifunctional protein 3) (Elongation factor p18) (Multisynthase complex auxiliary component p18)	
Immunogen	Synthesized peptide derived from human MCA3 AA range: 70-120	
Specificity	This antibody detects endogenous levels of human MCA3	
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.	
Source	Polyclonal, Rabbit,IgG	
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.	
Dilution	WB 1:1000-2000	
Concentration	1 mg/ml	
Purity	≥90%	
Storage Stability	-20°C/1 year	
Synonyms		
Synonyms Observed Band		
	Cytoplasm . Cytoplasm, cytosol . Nucleus . Cytoplasmic under growth arrest conditions. Translocated into the nucleus when growth resumes (S phase) and following DNA damage.	
Observed Band	conditions. Translocated into the nucleus when growth resumes (S phase) and	
Observed Band Cell Pathway	conditions. Translocated into the nucleus when growth resumes (S phase) and following DNA damage.	
Observed Band Cell Pathway Tissue Specificity	conditions. Translocated into the nucleus when growth resumes (S phase) and following DNA damage.	
Observed Band Cell Pathway Tissue Specificity Function	conditions. Translocated into the nucleus when growth resumes (S phase) and following DNA damage.	
Observed Band Cell Pathway Tissue Specificity Function Background matters needing	conditions. Translocated into the nucleus when growth resumes (S phase) and following DNA damage. Down-regulated in various cancer tissues.	



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Products Images			