



# PARVG rabbit pAb

<b>Catalog No</b>	YP-Ab-09869
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	PARVG
<b>Protein Name</b>	Gamma-parvin
<b>Immunogen</b>	Synthesized peptide derived from human PARVG AA range: 113-163
<b>Specificity</b>	This antibody detects endogenous levels of human PARVG
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cell junction, focal adhesion. Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cytoplasm, cytoskeleton . Constituent of focal adhesions. .
<b>Tissue Specificity</b>	Expressed predominantly in lymphoid organs, including spleen, thymus, lymph node, bone marrow and peripheral blood leukocytes and moderately in the digestive tract, including stomach, duodenum, jejunum, ileum, ileocecum and appendix, as well as in lung and liver. Also expressed in tumors, but at a lower level than in the corresponding normal tissues.
<b>Function</b>	
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



## Products Images