



# CN166 rabbit pAb

<b>Catalog No</b>	YP-Ab-09896
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	C14orf166 CGI-99
<b>Protein Name</b>	UPF0568 protein C14orf166 (CLE7 homolog) (CLE)
<b>Immunogen</b>	Synthesized peptide derived from human CN166 AA range: 29-79
<b>Specificity</b>	This antibody detects endogenous levels of human CN166
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Nucleus . Cytoplasm, cytosol . Cytoplasm, perinuclear region . Cytoplasm, cytoskeleton, microtubule organizing center, centrosome . May localize at the centrosome during mitosis (PubMed:15147888). Shuttles between the cytosol and the nucleus: enters into the nucleus in case of active transcription while it accumulates in cytosol when transcription level is low (PubMed:24608264). . ; Nucleus . Cytoplasm . (Microbial infection) Following influenza A virus (IAV) infection, included in influenza A virions via its association with packaged viral ribonucleoproteins (vRNP) in the nucleus and cytoplasm (PubMed:21900157, PubMed:26864902). .
<b>Tissue Specificity</b>	Widely expressed. Expressed at high level in heart and skeletal muscle. Expressed at intermediate level in liver, pancreas, fetal brain and fetal lung. Weakly expressed in adult brain, adult lung, placenta, fetal liver and fetal kidney. Overexpressed in many brain tumors.
<b>Function</b>	
<b>Background</b>	

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**