

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





CLCA1 rabbit pAb

Catalog No	YP-Ab-10130
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	CLCA1 CACC1
Protein Name	Calcium-activated chloride channel regulator 1 (Calcium-activated chloride channel family member 1) (hCLCA1) (Calcium-activated chloride channel protein 1) (CaCC-1) (hCaCC-1)
Immunogen	Synthesized peptide derived from human CLCA1 AA range: 192-242
Specificity	This antibody detects endogenous levels of human CLCA1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year Secreted, extracellular space . Cell membrane ; Peripheral membrane protein ; Extracellular side . Protein that remains attached to the plasma membrane
Storage Stability Synonyms Observed Band Cell Pathway	Secreted, extracellular space. Cell membrane; Peripheral membrane protein; Extracellular side. Protein that remains attached to the plasma membrane appeared to be predominantly localized to microvilli. Highly expressed in small intestine and colon namely in intestinal basal crypt epithelia and goblet cells, and appendix. Weakly expressed in uterus, testis and kidney. Expressed in the airways epithelium of both asthmatic and healthy patients. Expressed in the bronchial epithelium, especially in mucus-producing goblet cells. Expressed in normal turbinate mucosa and nasal polyp. Expressed
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	Secreted, extracellular space. Cell membrane; Peripheral membrane protein; Extracellular side. Protein that remains attached to the plasma membrane appeared to be predominantly localized to microvilli. Highly expressed in small intestine and colon namely in intestinal basal crypt epithelia and goblet cells, and appendix. Weakly expressed in uterus, testis and kidney. Expressed in the airways epithelium of both asthmatic and healthy patients. Expressed in the bronchial epithelium, especially in mucus-producing goblet cells. Expressed in normal turbinate mucosa and nasal polyp. Expressed
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	Secreted, extracellular space. Cell membrane; Peripheral membrane protein; Extracellular side. Protein that remains attached to the plasma membrane appeared to be predominantly localized to microvilli. Highly expressed in small intestine and colon namely in intestinal basal crypt epithelia and goblet cells, and appendix. Weakly expressed in uterus, testis and kidney. Expressed in the airways epithelium of both asthmatic and healthy patients. Expressed in the bronchial epithelium, especially in mucus-producing goblet cells. Expressed in normal turbinate mucosa and nasal polyp. Expressed



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images