

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com





GBRL1 rabbit pAb

Catalog No	YP-Ab-10135
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	GABARAPL1 GEC1
Protein Name	Gamma-aminobutyric acid receptor-associated protein-like 1 (Early estrogen-regulated protein) (GABA(A) receptor-associated protein-like 1) (Glandular epithelial cell protein 1) (GEC-1)
Immunogen	Synthesized peptide derived from human GBRL1 AA range: 32-82
Specificity	This antibody detects endogenous levels of human GBRL1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
	A wavefact
Concentration	1 mg/ml
Purity	1 mg/mi ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cytoplasmic vesicle, autophagosome . Cytoplasmic vesicle membrane ;
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasmic vesicle, autophagosome . Cytoplasmic vesicle membrane ; Lipid-anchor . Cytoplasm, cytoskeleton . Endoplasmic reticulum . Golgi apparatus . Ubiquitous. Expressed at very high levels in the brain, heart, peripheral blood leukocytes, liver, kidney, placenta and skeletal muscle. Expressed at very low levels in thymus and small intestine. In the brain, expression is particularly intense in motoneurons in the embryo and in neurons involved in somatomotor and neuroendocrine functions in the adult, particularly in the substantia nigra pars
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasmic vesicle, autophagosome . Cytoplasmic vesicle membrane ; Lipid-anchor . Cytoplasm, cytoskeleton . Endoplasmic reticulum . Golgi apparatus . Ubiquitous. Expressed at very high levels in the brain, heart, peripheral blood leukocytes, liver, kidney, placenta and skeletal muscle. Expressed at very low levels in thymus and small intestine. In the brain, expression is particularly intense in motoneurons in the embryo and in neurons involved in somatomotor and neuroendocrine functions in the adult, particularly in the substantia nigra pars
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasmic vesicle, autophagosome . Cytoplasmic vesicle membrane ; Lipid-anchor . Cytoplasm, cytoskeleton . Endoplasmic reticulum . Golgi apparatus . Ubiquitous. Expressed at very high levels in the brain, heart, peripheral blood leukocytes, liver, kidney, placenta and skeletal muscle. Expressed at very low levels in thymus and small intestine. In the brain, expression is particularly intense in motoneurons in the embryo and in neurons involved in somatomotor and neuroendocrine functions in the adult, particularly in the substantia nigra pars



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images