



# WIP12 Rabbit Polyclonal Antibody

<b>Catalog No</b>	YP-Ab-10267
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	IHC;IF
<b>Gene Name</b>	WIP12 CGI-50
<b>Protein Name</b>	WD repeat domain phosphoinositide-interacting protein 2 (WIP1-2) (WIP149-like protein 2)
<b>Immunogen</b>	Recombinant Protein of WIP12
<b>Specificity</b>	The antibody detects endogenous WIP12 protein
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	IHC-p 1:50-300. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	WD repeat domain phosphoinositide-interacting protein 2 (WIP1-2;WIP149-like protein 2)
<b>Observed Band</b>	49kD
<b>Cell Pathway</b>	Preautophagosomal structure membrane ; Peripheral membrane protein ; Cytoplasmic side . Localizes to omegasomes membranes which are endoplasmic reticulum connected structures at the origin of preautophagosomal structures. Enriched at preautophagosomal structure membranes in response to PtdIns3P. .
<b>Tissue Specificity</b>	Ubiquitously expressed (at protein level). Highly expressed in heart, skeletal muscle and pancreas. Expression is down-regulated in pancreatic and in kidney tumors.
<b>Function</b>	similarity:Contains 3 WD repeats.,tissue specificity:Ubiquitously expressed. Highly expressed in heart, skeletal muscle and pancreas. Expression is down-regulated in pancreatic and in kidney tumors.,
<b>Background</b>	WD repeat domain, phosphoinositide interacting 2(WIP12) Homo sapiens WD40 repeat proteins are key components of many essential biologic functions. They regulate the assembly of multiprotein complexes by presenting a beta-propeller platform for simultaneous and reversible protein-protein interactions. Members of the WIP1 subfamily of WD40 repeat proteins, such as WIP12, have a 7-bladed propeller structure and contain a conserved motif for interaction with phospholipids (Proikas-Cezanne et al., 2004 [PubMed 15602573]).[supplied by OMIM, Mar 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).