





BLK (Phospho-Tyr389) rabbit pAb

YP-Ab-10467
IgG
Human; Mouse;Rat
WB
BLK
BLK (Phospho-Tyr389)
Synthesized peptide derived from human BLK (Phospho-Tyr389)
This antibody detects endogenous levels of BLK (Phospho-Tyr389) at Human, Mouse,Rat
Liquid in PBS containing 50% glycerol, and 0.122% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
WB 1:500-2000
1 mg/ml
≥90%
-20°C/1 year
Tyrosine-protein kinase Blk (EC 2.7.10.2) (B lymphocyte kinase) (p55-Blk)
Cell membrane; Lipid-anchor. Present and active in lipid rafts. Membrane location is required for the phosphorylation of CD79A and CD79B (By similarity).
Expressed in lymphatic organs, pancreatic islets, Leydig cells, striate ducts of salivary glands and hair follicles.
catalytic activity:ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate.,function:May function in a signal transduction pathway that is restricted to B-lymphoid cells.,similarity:Belongs to the protein kinase superfamily. Tyr protein kinase family. SRC subfamily.,similarity:Contains 1 protein kinase
domain.,similarity:Contains 1 SH2 domain.,similarity:Contains 1 SH3 domain.,
domain.,similarity:Contains 1 SH2 domain.,similarity:Contains 1 SH3 domain., This gene encodes a nonreceptor tyrosine-kinase of the src family of proto-oncogenes that are typically involved in cell proliferation and differentiation. The protein has a role in B-cell receptor signaling and B-cell development. The protein also stimulates insulin synthesis and secretion in response to glucose and enhances the expression of several pancreatic beta-cell transcription factors. [provided by RefSeq, Aug 2010],



UpingBio technology Co.,Ltd

(Tel: 400-999-8863 **(** Emall:Upingbio.163.com



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images