



ROCK2 (Phospho-Tyr722) rabbit pAb

Catalog No	YP-Ab-10485
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	ROCK2 KIAA0619
Protein Name	ROCK2 (Phospho-Tyr722)
Immunogen	Synthesized peptide derived from human ROCK2 (Phospho-Tyr722)
Specificity	This antibody detects endogenous levels of ROCK2 (Phospho-Tyr722) at Human, Mouse, Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.141% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Rho-associated protein kinase 2 (EC 2.7.11.1) (Rho kinase 2) (Rho-associated, coiled-coil-containing protein kinase 2) (Rho-associated, coiled-coil-containing protein kinase II) (ROCK-II) (p164 ROCK-2)
Observed Band	
Cell Pathway	Cytoplasm. Cell membrane ; Peripheral membrane protein . Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasmic, and associated with actin microfilaments and the plasma membrane. .
Tissue Specificity	Expressed in the brain (at protein level).
Function	catalytic activity: ATP + a protein = ADP + a phosphoprotein. . enzyme regulation: Activated by RHOA binding. . function: Protein kinase that phosphorylates a large number of important signaling proteins, and thereby regulates the assembly of the actin cytoskeleton. Promotes formation of stress fibers and of focal adhesion complexes. Plays a role in smooth muscle contraction. . miscellaneous: Inhibited by Y-27632. . PTM: Phosphorylated upon DNA damage, probably by ATM or ATR. . similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. . similarity: Contains 1 AGC-kinase C-terminal domain. . similarity: Contains 1 PH domain. . similarity: Contains 1 phorbol-ester/DAG-type zinc finger. . similarity: Contains 1 protein kinase domain. . similarity: Contains 1 REM (Hr1) repeat. . subcellular location: Cytoplasmic, and associated with actin microfilaments and the plasma



membrane.,subunit:Homodim

Background

The protein encoded by this gene is a serine/threonine kinase that regulates cytokinesis, smooth muscle contraction, the formation of actin stress fibers and focal adhesions, and the activation of the c-fos serum response element. This protein, which is an isozyme of ROCK1 is a target for the small GTPase Rho. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of mouse spleen /Hela ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000