



EPH B1/3/4 (Phospho-Tyr778/792/774) rabbit pAb

Catalog No	YP-Ab-10497
Isotype	IgG
Reactivity	Human; Mouse; Rat
Applications	WB
Gene Name	EPHB1 ELK EPHT2 HEK6 NET
Protein Name	EPH B1/3/4 (Phospho-Tyr778/792/774)
Immunogen	Synthesized peptide derived from human EPH B1/3/4 (Phospho-Tyr778/792/774)
Specificity	This antibody detects endogenous levels of EPH B1/3/4 (Phospho-Tyr778/792/774) at Human, Mouse, Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.153% sodium azide.
Source	Polyclonal, Rabbit, IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Ephrin type-B receptor 1 (EC 2.7.10.1) (ELK) (EPH tyrosine kinase 2) (EPH-like kinase 6) (EK6) (hEK6) (Neuronally-expressed EPH-related tyrosine kinase) (NET) (Tyrosine-protein kinase receptor EPH-2)
Observed Band	
Cell Pathway	Cell membrane ; Single-pass type I membrane protein . Early endosome membrane . Cell projection, dendrite .
Tissue Specificity	Preferentially expressed in brain.
Function	catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., function: Receptor for members of the ephrin-B family. Binds to ephrin-B1, -B2 and -B3. May be involved in cell-cell interactions in the nervous system., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Ephrin receptor subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 1 SAM (sterile alpha motif) domain., similarity: Contains 2 fibronectin type-III domains., subunit: The ligand-activated form interacts with GRB2, GRB10 and NCK through their respective SH2 domains. The GRB10 SH2 domain binds EPHB1 through Tyr-928, while GRB2 binds residues within the catalytic domain. Interacts with EPHB6. The NCK SH2 domain binds EPHB1 through Tyr-594. Interacts with PRKCABP., tissue specificity: Preferentially expressed in brain.,

**Background**

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The protein encoded by this gene is a receptor for ephrin-B family members. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images