

**(** Tel: 400-999-8863 **(** Emall:Upingbio.163.com





## USF1 (Phospho-Thr153) rabbit pAb

| Catalog No         | YP-Ab-10517  |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human; Mouse;Rat   |
| Applications       | WB   |
| Gene Name          | USF1 BHLHB11 USF   |
| Protein Name       | USF1 (Phospho-Thr153)  |
| Immunogen          | Synthesized peptide derived from human USF1 (Phospho-Thr153)   |
| Specificity        | This antibody detects endogenous levels of USF1 (Phospho-Thr153) at Human, Mouse,Rat   |
| Formulation        | Liquid in PBS containing 50% glycerol, and 0.173% sodium azide.  |
| Source             | Polyclonal, Rabbit,IgG   |
| Purification       | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Dilution           | WB 1:500-2000  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | Upstream stimulatory factor 1 (Class B basic helix-loop-helix protein 11) (bHLHb11) (Major late transcription factor 1)  |
| Observed Band      |  |
| Cell Pathway       | Nucleus.   |
| Tissue Specificity | Kidney,  |
| Function           | disease:Genetic variations in USF1 are associated with combined hyperlipidemia type 1 (HYPLIP1) [MIM:602491]; also known as familial combined hyperlipidemia type 1 (FCHL1). HYPLIP1 is characterized by elevated levels of serum total cholesterol, triglycerides or both, and is observed in about 20% of individuals with premature coronary heart disease.,function:Transcription factor that binds to a symmetrical DNA sequence (E-boxes) (5'-CACGTG-3') that is found in a variety of viral and cellular promoters.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Binds DNA as an homodimer or a heterodimer (USF1/USF2). Interacts with varicella-zoster virus IE62 protein., |
| Background         | This gene encodes a member of the basic helix-loop-helix leucine zipper family, and can function as a cellular transcription factor. The encoded protein can activate transcription through pyrimidine-rich initiator (Inr) elements and E-box motifs. This gene has been linked to familial combined hyperlipidemia (FCHL).   |



## UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🗷 Emall:Upingbio.163.com



Alternative splicing of this gene results in multiple transcript variants. A related pseudogene has been defined on chromosome 21. [provided by RefSeq, Feb 2013],


matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western Blot analysis of 1 HeLa cell, 2 LPS 100ng/mL 30min treated ,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000