

📞 Tel: 400-999-8863 🗷 Email:Upingbio.163.com





# LIMK1 (Phospho-Ser310) rabbit pAb

Catalog No	YP-Ab-10548
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	LIMK1 LIMK
Protein Name	LIMK1 (Phospho-Ser310)
Immunogen	Synthesized peptide derived from human LIMK1 (Phospho-Ser310)
Specificity	This antibody detects endogenous levels of LIMK1 (Phospho-Ser310) at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.204% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	LIM domain kinase 1 (LIMK-1) (EC 2.7.11.1)
Observed Band	80kD
Cell Pathway	Cytoplasm . Nucleus . Cytoplasm, cytoskeleton . Cell projection, lamellipodium . Predominantly found in the cytoplasm. Localizes in the lamellipodium in a CDC42BPA, CDC42BPB and FAM89B/LRAP25-dependent manner
Tissue Specificity	Highest expression in both adult and fetal nervous system. Detected ubiquitously throughout the different regions of adult brain, with highest levels in the cerebral cortex. Expressed to a lesser extent in heart and skeletal muscle.
F 4!	catalytic activity (ATD La mateix — ADD La

**Function** 

catalytic activity:ATP + a protein = ADP + a phosphoprotein., disease:Haploinsufficiency of LIMK1 may be the cause of certain cardiovascular and musculo-skeletal abnormalities observed in Williams-Beuren syndrome (WBS), a rare developmental disorder de la 2014 and 1704 and syndrome involving genes from chromosome band 7q11.23.,function:Protein kinase which regulates actin filament dynamics. Phosphorylates and inactivates the actin binding/depolymerizing factor cofilin, thereby stabilizing the actin cytoskeleton. Isoform 3 has a dominant negative effect on actin cytoskeletal

changes. May be involved in brain development.,PTM:Autophosphorylated.,PTM:Phosphorylated on serine and/or threonine residues by ROCK1. May be dephosphorylated and inactivated by SSH1.,similarity:Belongs to the protein kinase superfamily. TKL Ser/Thr protein kinase family.,similarity:Contains 1 PDZ (DHR) doma



### UpingBio technology Co.,Ltd

📞 Tel: 400-999-8863 🗷 Email:Upingbio.163.com



#### Background

There are approximately 40 known eukaryotic LIM proteins, so named for the LIM domains they contain. LIM domains are highly conserved cysteine-rich structures containing 2 zinc fingers. Although zinc fingers usually function by binding to DNA or RNA, the LIM motif probably mediates protein-protein interactions. LIM kinase-1 and LIM kinase-2 belong to a small subfamily with a unique combination of 2 N-terminal LIM motifs and a C-terminal protein kinase domain. LIMK1 is a serine/threonine kinase that regulates actin polymerization via phosphorylation and inactivation of the actin binding factor cofilin. This protein is ubiquitously expressed during development and plays a role in many cellular processes associated with cytoskeletal structure. This protein also stimulates axon growth and may play a role in brain development. LIMK1 hemizygosity is implicated in the impaired visuospatial constructive cog


## matters needing attention

Avoid repeated freezing and thawing!

#### **Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

### **Products Images**



Western Blot analysis of 293T using primary antibody at 1:1000 dilution 4°C, overnight. Secondary antibody(catalog#:RS23920) was diluted at 1:10000 25°C, 1.5hours